arXiv:1210.5464v1 [cond-mat.soft] 19 Oct 2012

Hydrodynamic suppression of phase separation in active suspensions

Suzanne M. Fielding*
Department of Physics, University of Durham, Science Laboratories, South Road, Durham. DH1 3LE

(Dated: October 22, 2012)

We simulate a suspension of active squirming disks over the full range of volume fractions from dilute to close packed, with full hydrodynamics in two spatial dimensions. By doing so we show that "motility induced phase separation" (MIPS), recently proposed to arise generically in active matter, is strongly suppressed by hydrodynamic interactions. We give an argument for why this should be the case, and support it with counterpart simulations of active Brownian disks in a parameter regime more closely akin to hydrodynamic suspensions than in previous studies.

"Active matter" [1] comprises internal subunits that collectively drive the system far from Boltzmann equilibrium by each individually consuming energy. Biological examples include actively crosslinked polymeric filaments in the cell cytoskeleton [2]; cells grouped in living tissues [3]; suspensions of motile microorganisms [4, 5]; shoals of fish and flocks of birds [6]. Non-biological examples include vibrated granular monolayers [7–9], and self-propelled synthetic colloidal particles [10].

Distinct from the more familiar scenario of a passive complex fluid driven by (say) a global shear flow imposed at the system's boundaries, in active matter the driving out of equilibrium arises intrinsically in the active subunits throughout the system's own bulk. Consequently active materials can spontaneously develop mesoscopic or macroscopic mechanical stresses and deformations even without driving or loading from outside. Other exotic and generically emergent phenomena include swarming, pattern formation, giant number fluctuations, non-equilibrium ordering and phase separation. (For a recent review see Ref. [1].) These offer fascinating challenges to fluid dynamicist, rheologist and statistical physicist alike.

Many active particles are elongated and so have an intrinsic (steric) tendency to align with each other [11–15]. Activity mediated coupling between these orientational modes and fluctuations in the local number density then provides a generic mechanism for giant number fluctuations and phase separation [7, 9, 16], with the standard deviation ΔN in a subregion of material of mean number of particles N scaling as N^a with $a > 1/2$. (For passive systems away from any transition $a = 1/2$.)

Besides any such tendency for alignment, another generic mechanism for phase separation was recently put forward in the context of "run and tumble" particles, such as some species of motile bacteria [17, 18]. These swim in near straight-line runs at almost constant speed, between intermittent tumbles in which they suddenly randomize swim direction. The basic idea is that particles (a) accumulate where they move more slowly and (b) move more slowly where crowded. Positive (a)-(b) feedback gives rise to "motility-induced phase separation" (MIPS). This idea was recently extended analytically to active Brownian particles [19], consistent with simulations showing phase separation in active Brownian disks [20, 21]. These indeed lack any tendency for steric alignment; or for phase separation in equilibrium.

To date these simulations lack hydrodynamic interactions, in which moving particles set up flow fields that influence their neighbours. But such interactions arise widely in active matter [1], and this is fundamentally important because steady state properties in non-equilibrium systems depend strongly on dynamics (in contrast to equilibrium states, which depend only on the underlying free energy). This includes, crucially, the existence (or otherwise) of activity-induced phase separation.

The contribution of this work is to show that hydrodynamic interactions in fact suppress MIPS, which accordingly might not arise as generically in active matter as hitherto suggested, by simulating a suspension of active "squirming" disks [22–24], with full hydrodynamics, over the full range of particle volume fractions from dilute to close packed. (Being disks these lack any steric tendency to align, excluding a priori the first, non-MIPS mechanism for active phase separation discussed above. For simulations of active rods with hydrodynamics see Ref. [25]) We demonstrate carefully that hydrodynamics causes a key assumption of the (a) - (b) feedback mechanism of MIPS outlined above to fail. To support this, we further demonstrate suppression of MIPS in active Brownian disks in a parameter regime more closely akin to the squirmers, with MIPS recovered in the parameter regime explored previously for the Brownian particles [20].

Models — For both squirming and Brownian dynamics we simulate P disks of radius R in a box of area L^2. To prevent particle overlaps we impose a pairwise repulsive force $\boldsymbol{F}_{ij} = -f(a^3 - a^2)\hat{\boldsymbol{d}}_{ij}$, $a = 2sR/(d_{ij} - 2R)$ for interparticle separation $d_{ij} < 2R(1+s)$. The effective volume fraction is then $\phi = P\pi R^2(1+s)^2/L^2$.

The squirmers move through a Newtonian fluid of viscosity η by means of a prescribed tangential velocity $B_1 \sin\theta_i + B_2 \sin 2\theta_i$ round the disk edges. This gives a swim speed $v_0 = B_1/2$ [24] for a single disk undisturbed by any others, and an instantaneous swimming direction $\hat{\boldsymbol{e}}_i = (\cos\theta_i, \sin\theta_i)$ for the ith disk. These swim directions evolve over time due to hydrodynamic interactions between the disks, which are a priori unknown. Below we present data for $\beta \equiv B_2/B_1 = 0$, but have verified suppression of phase separation for $\beta = 0 \to \infty$.



FIG. 1: Active Brownian particles. Top: average particle speed as a function of average particle volume fraction. Expected spinodal shown by dotted line. Middle: scaled number fluctuations. Bottom: characteristic time $\tau_{\rm o}$ for reorientation of particle swim direction; and inter-particle collision time $\tau_{\rm c}$.


(a)$\zeta = 15.0, \phi = 0.5445$ (b)$\zeta = 15.0, \phi = 0.726$

(c)$\zeta = 1.0, \phi = 0.5445$ (d)$\zeta = 1.0, \phi = 0.726$

FIG. 2: Snapshots for active Brownian particles, corresponding to circles (a)-(d) in Fig. 1.

The active Brownian disks [20, 21, 26–28] have $\dot{\boldsymbol{r}}_i = v_0\hat{\boldsymbol{e}}_i + \mu\boldsymbol{F}_{ij}$, again with swim speed v_0 for a single undisturbed particle. In contrast to the squirmers their swim directions are unaffected by interparticle interactions, having independently prescribed dynamics $\dot{\theta}_i = n_i(t), \langle n_i(t)n_j(t')\rangle = 2\nu_r\delta_{ij}\delta(t-t')$, regardless of the frequency or closeness of interparticle encounters.

For both squirming and Brownian dynamics we simulate $P = 256$ particles, heavily constrained by computational cost for the squirmers (this P value being comparable with other squirmer studies [29]). A much larger P is achievable for the Brownian disks, but we present results only for $P = 256$ to ensure as direct a comparison as possible with the squirmers. We use units of length in which $L = 1$; time in which $v_0 = 1$; and mass in which $\eta = 1$ (squirmers) or $\mu = 1$ (Brownian). We set $f = 1, s = 0.1$ to ensure the dimensionless parameter $v_0/\mu f$ prescribing the small extent to which particles explore the interparticle repulsive potential is comparable to that in Ref. [20], giving almost hard disks.

The remaining dimensionless parameters are then the volume fraction $\phi = P\pi R^2(1+s)^2/L^2$; and for the Brownian disks also the ratio $\zeta = v_0\sqrt{P}/\nu_r L$ of the prescribed time of decorrelation of swim direction and the characteristic time interval between particle collisions.

Measured quantities— In what follows we report numerical results for the ensemble average particle swim speed $v = \frac{1}{PT}\int_0^T \sum_i |v_i(t)|$, averaged over the large time interval $t = 0 \to T$. (This speed differs slightly from the choice in Ref. [20], extracted from the early-time ballistic regime of the mean squared particle displacement, which one can show analytically is equivalent to $\left[\frac{1}{PT}\int_0^T \sum_i |v_i(t)|^2\right]^{1/2}$.) We also report the scaled number fluctuations $\delta_N = \Delta N/N^{1/2}$ in a region of the sample with average number of particles N.

We take the characteristic time $\tau_{\rm o}$ for decorrelation of particle swim direction to be the time interval $\Delta t = \tau_{\rm o}$ for the function $P(\Delta t) = \frac{1}{PT}\int_0^T dt \sum_i \boldsymbol{e}_i(t+\Delta t).\boldsymbol{e}_i(t)$ to fall to $1/e$. We find a reasonable measure of the time between inter-particle collisions to be $\tau_{\rm c} = \pi R/(v_0 - v_{\rm good})$ in which $v_{\rm good} = \frac{1}{PT}\int_0^T \sum_i \boldsymbol{v}_i(t).\boldsymbol{e}_i(t)$ is a measure of the extent to which a particle manages actually to achieve its full velocity in its attempted swim direction. The degree to which this differs from the free swim speed is a measure of the degree to which scattering occurs. We have checked by direct observation that $\tau_{\rm c}$ provides a reliable measure at low volume fraction. (At high volume fraction this comparison is more difficult to carry out, because particles slither round each other continuously.)

Results — We explore first active Brownian disks in the regime where the particle swim direction is slow to decorrelate, $\zeta \gg 1$. We do this to establish a point of contact with earlier simulation studies [20] of active Brownian disks, which were performed in this regime, to provide a context in which to discuss our new results below.

As seen in Fig. 1 (top), and correspondingly in Fig. 3 of Ref. [20] (though there with $P = 10^4$), the particle swim speed v decreases strongly with volume fraction ϕ. Accordingly this system is a good candidate for MIPS via the (a)-(b) feedback mechanism discussed above. The



FIG. 3: Hydrodynamic squirmers. Top: average particle speed as a function of average particle volume fraction. Expected spinodal shown by dotted line. Middle: scaled number fluctuations. Bottom: characteristic time τ_{o} for reorientation of particle swim direction; and inter-particle collision time τ_{c}.

expected spinodal, according to the instability criterion $d\log v/d\log\phi < -1$ of Ref. [17, 18], is located at the vertical dotted line in Fig. 1.

Phase separation is indeed observed: see the top two snapshots of Fig. 2, and Fig. 1 of Ref. [20]. This is reflected also in enhanced number fluctuations: our plot of $\delta_N(N)$ for each ϕ (not shown) shows all the same features as Fig. 3 of Ref. [28], though obviously cuts off sooner at high N due to our smaller number of simulated particles $P = 256$. Reporting in Fig. 1 (middle) the single value $\delta_{N=64}$, close to the peak of $\delta_N(N)$ for this P, clearly indicates separation with binodal onset around $\phi = 0.25$.

Consider now the hydrodynamic squirmers, our results for which are shown in Fig. 3. As for the Brownian disks the ensemble average swim speed declines strongly with particle volume fraction, suggesting that MIPS should again occur via the (a)-(b) feedback outlined above, with spinodal onset at the vertical dotted line according to the criterion $d\log v/d\log\phi < -1$ of Ref. [17]. Remarkably, however, we find no evidence for bulk phase separation in the squirmers: see the snapshots of Fig. 4 and the correspondingly suppressed number fluctuations $\delta_{N=64}$ in Fig. 3 (middle panel). (We do however find a tendency to form small string-like clusters at low volume fractions, as reported previously by other authors [29].)



(a)$\phi = 0.242$ (b)$\phi = 0.363$

(c)$\phi = 0.5445$ (d)$\phi = 0.7865$

FIG. 4: Snapshots for hydrodynamic squirmers, corresponding to circles (a)-(d) in Fig. 3.

Discussion — To understand how hydrodynamics might cause this suppression of motility-induced phase separation (MIPS) we now revisit the original argument for MIPS, as first put forward in the context of run and tumble dynamics and later generalised to other systems.

Model run and tumble particles move in a series of straight line runs at a constant speed v_0, when dilute, between tumbles in which they rapidly reassign their swim direction. These tumbles occur randomly with a typical intertumble time interval α^{-1}. The argument for phase separation stemming from this motility is as follows. (a) It can be shown that the local particle volume fraction in any small region of fluid scales inversely with the local swim speed v. (Although intuitively reasonable, this is in fact a strongly non-equilibrium effect stemming from activity. For passive colloidal particles the volume fraction is uniform in the absence of an external field.) (b) It is assumed that between tumbles particles move more slowly in regions of high volume fraction, being impeded by crowding, rendering the swim-speed a decreasing function of volume fraction $v(\phi) < v_0$. Positive feedback between (a) and (b) gives phase separation, onset with spinodal instability from an initially homogeneous state when $d\log v/d\log\phi < -1$. This argument extends to active Brownian particles of angular diffusivity ν_r, with an exact mapping $\alpha \to (d-1)\nu_r$ in d dimensions [19].

Consider carefully part (b) of this argument. In order meaningfully to define an intertumble swim speed $v(\phi) < v_0$ that is reduced by crowding, each particle must encounter many other particles between tumbles: the characteristic intertumble time α^{-1} must be large compared to the characteristic timescale τ_{c} between collisions. Transcribing this reasoning to Brownian particles, the angular diffusion time must be small compared to τ_{c}. Generalizing still further, we propose that regardless of the underlying microscopic dynamics – run and tumble, Brownian or hydrodynamic – the decay time $\tau_{\rm o}$ of the swim-direction autocorrelation function must greatly exceed the inter-particle collision timescale $\tau_{\rm c}$ if part (b) of the argument is to hold and MIPS is to occur.

With this reasoning in mind consider again Fig. 1 for Brownian disks in the regime $\zeta \gg 1$ discussed above. The bottom panel of this figure shows the decay time $\tau_{\rm o}$ of the autocorrelation function of particle swim direction, together with the typical time $\tau_{\rm c}$ between particle collisions. Consistent with the externally imposed angular diffusivity $\nu_{\rm r}$ being small in these runs (large $\zeta = 15$), $\tau_{\rm o}$ is a relatively large constant across all ϕ: except in the very dilute limit each particle encounters many others during the time it takes to alter its swim direction: $\tau_{\rm c} \ll \tau_{\rm o}$. Between angular reorientation events, then, each particle properly samples the ensemble average reduced swim speed $v(\phi) < v_0$: part (b) of the feedback loop holds and MIPS can indeed occur, as observed. Indeed, this mean-field nature of $v(\phi)$ was noted in Ref. [19].

In contrast, for the squirmers we are not at liberty to prescribe from the outset the timescale $\tau_{\rm o}$ of decorrelation of particle swim direction: instead this emerges naturally as a result of hydrodynamic interactions between the particles, which are a priori unknown. (Put differently, there is no externally tunable ζ for these particles.) Indeed, two-squirmer studies show that with hydrodynamics each scattering event ("collision") typically results in an $O(1)$ change in swim-direction for each particle involved. Accordingly, in these many-squirmer simulations we expect $\tau_{\rm o} \approx \tau_{\rm c}$. This is indeed observed: Fig. 3 (bottom panel). Squirmers will therefore be unable properly to sample the reduced $v(\phi) < v_0$ between reorientation events: part (b) of the feedback argument fails, and MIPS is suppressed.

To support this argument we finally revisit the Brownian disks, but now imposing a smaller $\zeta = 1$ so that the particles reorient their swim directions much more quickly than in the large ζ case discussed above, giving dynamics more closely akin to the squirmers ($\tau_{\rm o} \approx \tau_{\rm c}$, Fig. 1, bottom). Phase separation is then indeed strongly suppressed: see the bottom two snapshots of Fig. 2, and the significantly reduced number fluctuations of Fig. 1 (middle panel). This is despite the ensemble average velocity $v(\phi)$ still decreasing strongly enough for this $\zeta = 1$ to satisfy the condition for spinodal onset. Put simply, particles that rapidly reorient do not have time to sample this mean field $v(\phi)$ in between reorientation events.

Despite the absence of true bulk phase separation for the squirmers, and for the Brownian disks with $\tau_{\rm o} \approx \tau_{\rm c}$, some particle clustering is nonetheless still apparent in Figs. 2c,d and Fig. 4a-d. Whether this clustering can be interpreted in terms of a nearby but suppressed MIPS, which could therefore be a generic feature of active matter with hydrodynamics, remains an open question.

Conclusions — We have simulated a suspension of active squirming disks across the full range of volume fractions from dilute to close packed, with full hydrodynamics. In doing so we have shown that hydrodynamic interactions strongly suppress motility induced phase separation. These findings should apply generically to active systems in which hydrodynamics are important, and in which the effective particle velocity depends on volume fraction via collisions impeding particle motion. Obvious exceptions include systems in which v depends on ϕ instead by chemically mediated mechanisms (such as quorum sensing [30]), allowing MIPS to arise at relatively low volume fractions, and/or systems in which the particles reside in (or on the surface of) a gel [31]. Such exceptions notwithstanding, the mechanism proposed here is expected to arise widely in active matter, and particularly in active colloids, which form the focus of intense current experimental interest [10].

Acknowledgements The author thanks Mike Cates, Paul Chaikin and Oliver Harlen for discussions. The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC Grant agreement no. 279365

---

[*] Electronic address: suzanne.fielding@durham.ac.uk

[1] M. C. Marchetti *et al.*, arXiv:1207.2929 and references therein (2012).

[2] F. Jülicher, K. Kruse, J. Prost, and J.-F. Joanny, Physics Reports **449**, 3 (2007).

[3] M. Poujade *et al.*, Proc. Nat. Acad. Sci. U.S.A. **104**, 15988 (2007).

[4] H. C. Berg, *E. coli in Motion* (Springer, New York, 2003).

[5] S. Rafai, L. Jibuti, and P. Peyla, Phys. Rev. Lett. **104**, 098102 (2010).

[6] J. K. Parrish and W. M. Hamner, *Three Dimensional Animal Groups* (Cambridge University Press, Cambridge, England, 1997).

[7] V. Narayan, S. Ramaswamy, and N. Menon, Science **317**, 105 (2007).

[8] D. V. A. Kudrolli, G. Lumay and L. S. Tsimring, Phys. Rev. Lett. **100**, 058001 (2008).

[9] J. Deseigne, O. Dauchot, and H. Chate, Phys. Rev. Lett. **105**, 098001 (2010).

[10] G. Ruckner and R. Kapral, Phys. Rev. Lett. **98**, 150603 (2007); J. Howse*et al.*, Phys. Rev. Lett. **99**, 048102 (2007); J. Palacci, C. Cottin-Bizonne, C. Ybert, and L. Bocquet, Phys. Rev. Lett. **105**, 088304 (2010); A. Erbe *et al.*, J. Phys. - Cond. Matt. **20**, (2008); W. Paxton *et al.*, J. Am. Chem. Soc. **126**, 13424 (2004); Y. Hong *et al.*, Phys. Rev. Lett. **99**, 178103 (2007); H.-R. Jiang, N. Yoshinaga, and M. Sano, Phys. Rev. Lett. **105**, 268302 (2010); G. Volpe *et al.*, Soft Matter **7**, 8810 (2011); I. Theurkauff *et al.*, Phys. Rev. Lett. **108**, 268303 (2012).

[11] J. Toner and Y. Tu, Phys. Rev. E **58**, 4828 (1998).

[12] J. Toner and Y. Tu, Phys. Rev. Lett. **75**, 4326 (1995).

[13] S. Ramaswamy, R. Simha, and J. Toner, Europhys. Lett. **62**, 196 (2003).

[14] J. Toner, Y. Tu, and S. Ramaswamy, Annals Of Physics **318**, 170 (2005).

[15] A. Baskaran and M. C. Marchetti, Phys. Rev. E **77**, 011920 (2008); A. Baskaran and M. C. Marchetti, Phys. Rev. Lett. **101**, 268101 (2008).

[16] H. Chate, F. Ginelli, and R. Montagne, Phys. Rev. Lett. **96**, 180602 (2006); H. Chate *et al.*, Eur. Phys. J. B **64**, 451 (2008).

[17] J. Tailleur and M. E. Cates, Phys. Rev. Lett. **100**, 218103 (2008).

[18] M. E. Cates, Rep. Prog. Phys. **75**, 042601 (2012).

[19] M. E. Cates and J. Tailleur, arXiv:1206.1805 (2012).

[20] Y. Fily and M. C. Marchetti, Phys. Rev. Lett. **108**, 235702 (2012).

[21] G. S. Redner, M. F. Hagan, and A. Baskaran, arXiv:1207.1737v3 (2012).

[22] M. J. Lighthill, Comm. Pure Appl. Math. **5**, 109 (1952).

[23] J. R. Blake, J. Fluid Mech. **46**, 199 (1971).

[24] J. R. Blake, Bulletin of the Australian Mathematical Society **5**, 255 (1971).

[25] D. Saintillan and M. J. Shelley, Phys. Rev. Lett. 99, 058102 (2007); ibid. 100, 178103 (2008).

[26] J. Bialke, T. Speck, and H. Löwen, Phys. Rev. Lett. **108**, 168301 (2012).

[27] S. R. McCandlish, A. Baskaran, and M. F. Hagen, Soft Matter **8**, 2527 (2012).

[28] S. Henkes, Y. Fily, and M. C. Marchetti, Phys. Rev. E **84**, 040301 (2011).

[29] T. Ishikawa, J. Locsei, and T. Pedley, J. Fluid Mech. **615**, 401 (2008).

[30] M. B. Miller and B. L. Bassler, Ann. Rev. Microbiol. **55**, 165 (2001).

[31] M. Cates, D. Marenduzzo, I. Pagonabarraga, and J. Tailleur, Proc. Nat. Acad. Sci. U. S. A. **107**, 11715 (2010); F. D. C. Farrell, M. C. Marchetti, D. Marenduzzo, and J. Tailleur, Phys. Rev. Lett. **108**, 248101 (2012); C. L. Liu *et al.*, Science **334**, 238 (2011); X. F. Fu *et al.*, Phys. Rev. Lett. **108**, 198102 (2012).